

De Novo Mutation of the β -Globin Gene Initiation Codon (ATG→AAG) in a Northern European Boy

John S. Waye,^{1,2*} Barry Eng,¹ Margaret Patterson,¹ Ronald D. Barr,³ and David H.K. Chui^{1,2}

¹Provincial Hemoglobinopathy DNA Diagnostic Laboratory, McMaster University Medical Centre, Hamilton, Ontario, Canada

²Department of Pathology, McMaster University School of Medicine, Hamilton, Ontario, Canada

³Department of Pediatrics, McMaster University School of Medicine, Hamilton, Ontario, Canada

We present a case of β -thalassemia intermedia involving a 13-year-old boy of Northern European descent. His mother, father and older sister have normal hematologic indices. Molecular studies demonstrate that the proband carries a novel mutation of the β -globin gene initiation codon (ATG→AAG) which should give rise to β^0 -thalassemia trait. The possibility of non-paternity was excluded, indicating that the novel mutation was the result of a de novo event. A review of the literature indicates that mutations involving the β -globin gene initiation codon can give rise to a more severe phenotype than is generally associated with most other β^+ or β^0 mutations. *Am. J. Hematol.* 56:179–182, 1997.

© 1997 Wiley-Liss, Inc.

Key words: β -thalassemia; initiation codon; de novo mutation

INTRODUCTION

β -thalassemia is one of the most common single-gene disorders of man, occurring at high frequencies among individuals of Mediterranean, East Indian, Middle Eastern, African, or Southeast Asian descent [1]. More than 150 different β -thalassemia mutations have been reported, most of which are single base substitutions or small insertions or deletions within the β -globin gene [2].

The phenotypic severity of individual β -thalassemia mutations can vary considerably depending on the molecular defect [3–5]. The mildest mutations, designated β^+ , are those that reduce the expression of normal β -chains. β^0 mutations abolish expression of normal β -chains and therefore are more severe than β^+ mutations. The most severe class of mutations is the so-called inclusion body or dominant β -thalassemia mutations that give rise to highly unstable β -chain variants [6–8]. Carriers of such mutations exhibit a relatively severe β -thalassemia intermedia phenotype.

In this report, we describe a case of β -thalassemia intermedia due to a de novo mutation of the β -globin gene initiation codon. A review of the literature confirms that this type of mutation gives rise to a more severe

TABLE I. Hematology Results for the Family Under Study

	Father (48 years)	Mother (42 years)	Proband (13 years)	Sister (15 years)
Hb (g/dL)	14.8	12.7	9.0	12.6
Hct	0.424	0.360	0.270	0.356
RBC ($10^{12}/L$)	4.58	3.95	4.87	4.29
MCV (fL)	92.4	92.0	55.0	83.0
MCH (pg)	32.2	32.0	18.6	29.4
RDW	13.7	13.5	26.9	12.7
Hb A (%)	96.1	95.3	87.9	95.7
Hb A ₂ (%)	3.3	3.0	5.7	3.1
Hb F (%)	0.6	1.7	6.4	1.2
α -genotype	$\alpha\alpha/\alpha\alpha$	$\alpha\alpha/\alpha\alpha$	$\alpha\alpha/\alpha\alpha$	$\alpha\alpha/\alpha\alpha$

phenotype than is generally associated with most other β^+ or β^0 thalassemia mutations.

Contract grant sponsor: Ontario Thalassemia Foundation.

*Correspondence to: John S. Waye, Ph.D., Department of Pathology, Room 3N17, McMaster University Medical Centre, 1200 Main Street West, Hamilton, Ontario, Canada L8N 3Z5.

Received for publication 8 November 1996; Accepted 11 June 1997

TABLE II. Hematologic Phenotypes of Heterozygotes for β -Globin Gene Initiation Codon Mutations

Mutation	Population	Reference	Sex-age	Hb (g/dL)	MCV (fL)	Hb A ₂ (%)	Hb F (%)
ATG → GTG	Japanese	Hattori et al. [12]	M-13	11.6	53.0	6.7	1.9
			F-41	9.9	56.0	6.0	0.8
			F-51	11.0	55.0	6.2	1.5
			F-54	9.1	70.0	6.2	7.1
ATG→ACG	Swiss	Beris et al. [14]	M	11.5	59.0	4.0	3.2
			F	9.4	60.0	7.3	3.1
	Yugoslavian	Jankovic et al. [15]	M-52	11.5	57.6	6.2	0.6
			M-25	11.6	54.1	5.9	0.9
			F-23	10.0	56.1	6.0	0.7
			M-25	11.6	54.1	5.9	0.9
	Belgian	Wildmann et al. [16]	F-70	10.2	55.4	5.9	1.1
			M-37	11.8	56.1	5.4	0.9
			M-34	12.5	54.0	6.1	1.1
			F-11	10.2	50.7	6.3	0.5
			M-07	10.0	53.3	6.4	1.8
			F-06	9.2	50.1	5.4	2.3
			F-05	9.4	53.4	5.9	4.4
			M-04	9.9	48.5	5.8	6.8
			F-01	9.1	51.4	4.1	22.0
ATG→AGG	Korean	Koo et al. [18]	F-53	10.4	53.0	6.1	2.6
			F-50	10.6	58.0	6.4	2.2
			F-48	11.0	62.0	5.7	4.7
			M-25	12.9	52.0	6.1	0.7
			M-16	11.3	49.0	6.6	3.8
			F-25	10.5	76.0	3.9	21.7
			F-15	9.6	67.0	4.8	11.8
			M-13	11.0	61.0	5.7	16.0
ATG→ATA	Swedish	Landin et al. [19]	M-73	10.2	54.8	5.9	0.6
			F-47	10.7	55.2	5.2	<0.5
			M-35	12.0	53.4	5.9	<0.5
			M-21	12.1	56.4	4.7	<0.5
			F-17	9.6	56.1	5.0	1.2
			M-2	9.5	48.9	5.5	4.4
			M-1	9.6	49.1	5.5	4.4
ATG→ATT	U.S. Caucasian	Rahbar and Nozari [21]	F-23	7.7	74.0	6.4	2.0
ATG→AAG	N. European	This report	M-13	9.0	55.0	5.7	6.4

TABLE III. Hematologic Phenotypes of Heterozygotes for Different β -Thalassemia Mutations*

Mutation	Severity	Hb (g/dL)	MCV (fL)	Hb A ₂ (%)
-87 C→G	Mild β^+	14.1 ± 1.4 (male)	74.0 ± 5.2	5.3 ± 0.7
IVS-I-6				
T→C	Mild β^+	13.8 ± 0.7 (male)	70.4 ± 1.9	3.9 ± 0.4
IVS-I-110				
G→A	Severe β^+	13.2 ± 1.0 (male)	63.8 ± 2.7	4.5 ± 0.5
IVS-II-745				
C→G	Severe β^+	13.1 ± 0.5 (male)	62.5 ± 3.0	5.0 ± 0.5
IVS-I-1				
G→A	Severe β^0	12.8 ± 0.8 (male)	63.3 ± 3.1	4.8 ± 0.4
Codon 39				
C→T	Severe β^0	13.3 ± 0.9 (male)	62.9 ± 3.0	5.3 ± 0.7
Initiation	Severe β^0	11.1 ± 1.1 (male)	56.4 ± 6.6	5.7 ± 0.7
codon		9.9 ± 0.8 (female)		

*The values given for initiation codon mutations (mean ± SD) were derived from Table II. The values given for all other mutations were reported by Rosatelli et al. [22].




Fig. 1. Nucleotide sequence analysis of PCR amplified β -globin gene sequences from the proband. The normal initiation codon (ATG) is boxed.

CASE REPORT

The proband is a 13-year-old boy of Northern European descent who presented with the hematologic profile of β -thalassemia intermedia (Hb = 9.0 g/dL, MCV = 55.0 fL, Hb A₂ = 5.7%, Hb F = 6.4%). The blood film showed hypochromia, microcytosis, targeting, and stippling. The physical examination revealed splenomegaly, palpable at 1 cm below the left costal margin. The proband's parents and older sister have normal hematologic indices (Table I).

Molecular studies were initiated on the proband. The β -globin gene was amplified using the polymerase chain reaction (PCR) [9] and the PCR product was sequenced using the dideoxy termination method [10]. As shown in Figure 1, the proband is heterozygous for a single base substitution involving the initiation codon (ATG \rightarrow AAG). This mutation abolishes the recognition site for the restriction endonuclease *Nco* I. PCR amplified DNA from the family members was digested with *Nco* I and analysed by polyacrylamide gel electrophoresis. The results demonstrate that only the proband carries the initiation codon mutation. The possibility of non-paternity

was investigated by RFLP typing of VNTR loci D2S44 and D5S110 and by PCR typing of the HLA-DQ α locus [11] (data not shown). The test results were fully consistent with paternity as stated, with a paternity index of 3,888 and a probability of paternity in excess of 99.97% (prior probability = 0.5). Therefore, the novel initiation codon mutation was the result of a de novo event.

DISCUSSION

Mutations of the β -globin gene initiation codon have been described in several different populations. Including the novel ATG \rightarrow AAG mutation presented in this report, seven of the nine possible single-base substitutions have been reported [12–22]. It is of interest that the hematologic profile of the patient described in this report is more severe than typical β -thalassemia trait. In particular, the proband's microcytic anemia (Hb = 9.0 g/dL, MCV = 55.0 fL) is quite striking for someone who carries only one β -thalassemia allele.

As noted by Landin et al. [19], severe microcytic anemia appears to be a common feature associated with mutations of the initiation codon of the β -globin gene. A review of 35 patients from the literature, all of whom are carriers of mutations involving the initiation codon of the β -globin gene, reveals that both the hemoglobin level and MCV are markedly reduced relative to typical β -thalassemia trait (Tables II and III) [23,24]. It is thought that mutations affecting mRNA translation are associated with decreased cytosolic mRNA levels, most likely due to an intranuclear event or the nuclear-cytoplasm transport process [25,26]. In the case of initiation codon mutations, this effect may be exaggerated and thereby account for the severely abnormal hematologic indices. Another possibility may be that translation of the mutant mRNA initiates from the single in-frame initiation codon (codon 55) and produces a truncated β -chain that is highly unstable. Proteolytic degradation of the truncated β -chains would allow free α -chains to accumulate and precipitate, possibly resulting in ineffective erythropoiesis and/or peripheral hemolysis. In this manner, initiation codon mutations may resemble the exon III mutations that are associated with dominant β -thalassemia [6–8].

ACKNOWLEDGMENTS

We sincerely thank Dr. W. Nicholson for referring this case to our attention, and the family for their cooperation. This work was funded in part through a research grant from the Ontario Thalassemia Foundation (J.S.W.).

REFERENCES

1. Weatherall DJ, Clegg JB: "The Thalassemia Syndromes," 3rd ed. Oxford: Blackwell Scientific Publications, 1981.

2. Baysal E, Carver MF: The β - and δ -thalassemia repository, 8th ed. Hemoglobin 19:213, 1995.
3. Kazazian HH Jr: The thalassemia syndromes: Molecular basis and prenatal diagnosis in 1990. Semin Hematol 27: 209, 1990.
4. Loukopoulos D: Thalassemia: Genotypes and phenotypes. Ann Hematol 62:85, 1991.
5. Cao A, Galanello R, Rosatelli MC: Genotype-phenotype in β -thalassemia. Blood Rev 8:1, 1994.
6. Thein SL: Dominant β -thalassaemia: Molecular basis and pathophysiology. Br J Haematol 80:272, 1992.
7. Thein SL, Hesketh C, Taylor P, Temperley IJ, Hutchinson RM, Old JM, Wood WG, Clegg JB, Weatherall DJ: Molecular basis for dominantly inherited inclusion body β -thalassemia. Proc Natl Acad Sci USA 87:3924, 1990.
8. Kazazian HH Jr, Dowling CE, Hurwitz RL, Coleman M, Stopeck A, Adams JG III: Dominant thalassemia-like phenotypes associated with mutations in exon 3 of the β -globin gene. Blood 79:3014, 1992.
9. Saiki RK, Gelfand DH, Stoffel S, Scharf SJ, Higuchi R, Horn GT, Mullis KB, Erlich HA: Primer-directed enzymatic amplification of DNA with a thermostable polymerase. Science 239:487, 1988.
10. Sanger F, Nicklen S, Coulson AR: DNA sequencing with chain termination inhibitors. Proc Natl Acad Sci USA 74:5463, 1977.
11. Waye JS: Forensic identity testing using highly polymorphic DNA markers: Current status and emerging technologies. Transf Med Rev 7:193, 1993.
12. Hattori Y, Yamashiro Y, Ohba Y, Miyaji T, Morishita M, Yamamoto Ku, Yamamoto Ki, Narai S, Kimura A: A new β -thalassemia mutation (initiation codon ATG \rightarrow GTG) found in the Japanese population. Hemoglobin 15:317, 1991.
13. Ringelhann B, Szelenyi JG, Horanyi M, Svobodova M, Divoky V, Indrak K, Hollán S, Marosi A, Laub M, Huisman THJ: Molecular characterization of β -thalassemia in Hungary. Hum Genet 92:385, 1993.
14. Beris P, Darbellay R, Speiser D, Kirchner V, Miescher PA: De novo initiation codon mutation (ATG \rightarrow ACG) of the β -globin gene causing β^0 -thalassemia in a Swiss family. Am J Hematol 42:248, 1993.
15. Jankovic L, Efremov GD, Josifovska O, Juricic D, Stoming TA, Kutlar A, Huisman THJ: An initiation codon mutation as a cause of a β -thalassemia. Hemoglobin 14:169, 1990.
16. Wildmann C, Larondelle Y, Vaerman JL, Eeckels R, Martiat P, Philippe M: An initiation codon mutation as a cause of β -thalassemia in a Belgian family. Hemoglobin 17:19, 1993.
17. Lam VMS, Xie SS, Tam JWO, Gu YL, Li AMC: A new single nucleotide change at the initiation codon (ATG \rightarrow AGG) identified in amplified genomic DNA of a Chinese β -thalassemic patient. Blood 75:1207, 1990.
18. Koo MS, Kim SI, Cho HI, Hattori Y, Yamashiro Y, Hoshitani M, Ohba Y, Miyaji T, Yamamoto Ku, Yamamoto Ki: A β -thalassemia mutation found in Korea. Hemoglobin 16:313, 1992.
19. Landin B, Rudolphi O, Ek B: Initiation codon mutation (ATG \rightarrow ATA) of the β -globin gene causing β -thalassemia in a Swedish family. Am J Hematol 48:158, 1995.
20. Saba L, Meloni A, Sardu R, Travi M, Primignani P, Rosatelli MC, Cao A: A novel β -thalassemia mutation (G \rightarrow A) at the initiation codon of the β -globin gene. Hum Mutat 1:420, 1992.
21. Rahbar S, Nozari G: A novel initiation codon mutation (ATG \rightarrow ATT) in a β -thalassemia patient. Hemoglobin 17:557, 1993.
22. Ohba Y, Hattori Y, Harano T, Harano K, Fukumaki Y, Ideguchi H: β -Thalassemia mutations in Japanese and Koreans. Hemoglobin 21:191, 1997.
23. Rosatelli C, Leoni GB, Tuveri T, Scalas MT, Mosca A, Galanello R, Gasperini D, Cao A: Heterozygous β -thalassemia: Relationship between the hematological phenotype and the type of β -thalassemia mutation. Am J Hematol 39:1, 1992.
24. Rund D, Fillon D, Strauss N, Rachmilewitz EA, Oppenheim A: Mean corpuscular volume of heterozygotes for β -thalassemia correlates with the severity of the mutations. Blood 79:238, 1992.
25. Takeshita K, Forget BG, Scarpa A, Benz EJ Jr: Intranuclear defect in β -globin mRNA accumulation due to premature translation termination codon. Blood 64:13, 1984.
26. Humphries RK, Ley TJ, Anagnou NP, Baur AW, Nienhuis AW: β^0 -39 Thalassemia gene: A premature termination codon causes β -mRNA deficiency without affecting cytoplasmic β -mRNA stability. Blood 64:23, 1984.
27. Baserga SJ, Benz EJ Jr: β -Globin nonsense mutation: deficient accumulation of mRNA occurs despite normal cytoplasmic stability. Proc Natl Acad Sci USA 89:2935, 1992.